Code No. 2965/AICTE/M

FACULTY OF ENGINEERING

B.E. (CSE/EEE/Inst.) (AICTE) IV-Semester (Main) Examination, December 2020

Subject : Mathematics - III (P & S)

Time: 2 Hours

Max. Marks: 70

Note: (Missing data if, any can be assumed suitable).

PART - A

Answer any five questions.

(5 x 2 = 10 Marks)

1 Define conditional probability.

2 A continuous random variate X has the probability density function

f(x) = a + bx, $0 \le x \le 1$

= 0, elsewhere of the distribution is $\frac{1}{2}$, find the values of a and b.

- 3 Define Binomial distribution.
- 4 Define Skewness
- 5 Find the mean of Exponential distribution.
- 6 Explain Normal distribution
- 7 Write normal equations of straight line.
- 8. Write the equations of the regression lines.
- 9 Define Null hypothesis.
- 10 Define Population and sample.

PART - B

Answer any four questions.

(4 x 15 = 60 Marks)

- 11 (a) State and prove theorem of total probability.
 - (b) If A and B are two mutually exclusive events of a random experiment, then P(A U B) = P(A) + P(B)
- 12 (a) If the probability of a bad reaction from a certain injection is 0.001, determine the chance that out to 2.000 individuals more than 2 will get bad reaction.
 - (b) Calculate the quartile coefficient of skewness from the following data.

Weight (this)	70-80	80-90	90-100	100-110	110-120	120-130	130-140	140-150
Weight (lbs) No. of persons	12	18	35	42	50	45	20	8

13 (a) A continuous random variable X has probability density function

$$\digamma(\pi)=\frac{1}{4}(x^3+1),\ 0\leq\pi\leq1.\ \text{Find 'a' such that }P(X\leq a)\equiv P(X\geq a),$$

- (b) A continuous random variable X is uniformly distributed with mean 1 and variance 3. Find P(X < 0).</p>
- 14 (a) A coin was tossed 400 times and head turned up 216 times. Test the hypothesis that the coin is unbiased at 5% level of significance.
 - (b) Fit a parabola y = a + bx + cx2 for the following data:

X	1	2	3	4	5	6
f(x)	1000	800	500	300	150	90

15 (a) Five dice were thrown 96 times and the no. of times 4, 5 or 6 were thrown were:

No. of dice showing 4, 5, or 6	5	4	3	2	1	0
Frequency	8	18	35	24	10	1

Find the probability of getting this result by chance.

- (b) A normal population has mean 0. 1 and a S.D. of 2.1. Find the probability that mean of simple sample of 900 members will be negative?
- 16 (a) The probability density function of variate X is

X	0	1	2	3	4	5	-6
P(X)	K	3K	5K	7K	9K	11K	13K

Find P(X < 4), $P(X \ge 5)$, $P(3 < X \le 6)$

- (b) If X is a Poisson variate such that P(X = 2) = 3P(X = 4) + 45P(X = 6). Find the mean and variance of X.
- 17 (a) Find the moment generating function of uniform distribution.
 - (b) Find the correlation coefficient and regression lines for the following data:

х	1	2	3	4 8	5
y.	2	5	3	8	7

FACULTY OF ENGINEERING

B.E. IV - Semester (EE/Inst/CSE) (AICTE) (Main&Backlog) Examination,

October 2021

Subject: Mathematics - III (Probability and Statistics)

Time: 2 Nours

Max. Marks: 70

(Missing data, if any, may be suitably assumed)

PART - A

Note: Answer any five questions.

(5x2 = 10 Marks)

State Baye's theorem.

the following probability distribution: 2

Mr.	14	1.2	3	4
Α:	1		2140	215
P(X):	1/10	1/5	3/10	Alst

3 Average number of accidents on any day on a national highways 1.8. Find the probability that the number of accidents is at least one. 5/817

4 Define Skewness of a distribution.

5 Write any two properties of the normal probability curve

7 Two lines of regression are 7x-16y+9 = 0, 5y=1x-3=0. Find \$\bar{x}\$ and \$\bar{y}\$.
 8 Define type I and type II errors.

9 Write the assumptions for conducting t-tests

10 Explain briefly F-test.

PART-B

Note: Answer any four questions

(4x15 = 60 Marks)

11 (a) State and prove theorem of total probability.

(b) A continuous random variable X has the pdf $f(x) = \begin{cases} ax^3, 0 \le x \le 1 \\ 0 & olsowhere \end{cases}$

Find (i) a (ii) $P\left(X < \frac{1}{4}\right)$ and (iii) $P\left(X > \frac{1}{2}\right)$.

12 (a) Fit a binomial distribution to the following data:

T !	To	11	2	3	4	5
1:	2	14	20	34	22	8

(b) Calculate the first four moments about the mean for the following data:

r. 1	2	13	4	5	0	7	В	9
6	- 6	13	25	30	22	9	1.5	1.2

- 13 (a) The marks obtained in Mathematics by the students in a class are approximately normally distributed with mean 62 and variance 36. If 3 students are selected at random, find the probability that at least one of them would score more than 80 marks.
 - (b) Find the mean and moment generating function of exponential distribution.

14 (a) Find the least square line y = a + for for the following data:

x: 1-2	1-7	0	13:	12	20 50 50
1 1	2	3	3	4	7,75

(b) Intelligence test of two groups of boys and girls gave the following results.

Girls: 1,=84, S.D. = 10, M,=121

Boys: 1, =80, S.D. = 14, n, = 81.

is the difference between the standard deviations significant? Test at 5% level of significance.

15 The values of two random samples are given below.

Sample A 15 25 16 20 22 24 21 17 19 23
Sample B 35 31 25 38 26 29 32 34 33 27 29 31
Can we conclude that the two samples are drawn from the same copulation? Test at 55 level of significance.

16 (a) A dice is thrown twice and the sum of Lie numbers appearing is noted to be 8. Find the probability that the number 5 has appeared at least once.

(b) Find the variance of the normal distribution.

17 From the following data, calculate coefficient of correlation between X and Y and the two lines of regression equations.

Χ;		12	3	4 6/5	6	7	8	9
Y:	12	111	13	15 14	157	18	10	10

Code No: 15050/AICTE/BI

FACULTY OF ENGINEERING

BE III - Semester (AICTE) (CME/IT) (Backlog) Examination, November 2021

Subject: Mathematics-III (Probability & Statistics)

Time: 2 Hours

Max .Marks: 70

Note: Missing data, if any, may be suitably assumed

PART - A

Answer any five questions.

(5x2=10 Marks)

1 State Baye's theorem.

2 A random variable X has the following probability distribution:

Y	0	1	2	3
P(x)		1	3.	104
SI APOLA	6	3	10	30

3. A die is thrown 4 times. Getting a number greater than 2 is a success. Find the probability of getting exactly one success.

4 Find the moment generating function of poisson distribution.

5. A continuous random variable X is uniformly distributed with mean 1 and variance 3. Find P(X<0)

6 Define exponential distribution.

7 Write normal equations for fitting a straight-line of the form y = a + bx.

8 Show that the coefficient of correlation is the geometric mean of the regression coefficients.

9 Explain briefly F-test.

10 Write any two applications of Chu-square test.

PART - B

Answer any four questions.

(4x15= 60 Marks)

11 (a) State and prove theorem of total probability.

- (b) A factory has four independent units A, B, C and D which produce 40%, 30%, 20% and 10% of identical items, respectively. The percentages of defective items produced by these units are 2%, 1%, 0.5% and 0.25% respectively. If an item is selected at random, find the probability that the item is defective.
- 12 (a) Find the mean and variance of binomial distribution.
 - (b) If X is a poisson variate such that $3P(x=4)=\frac{1}{3}P(x=2)+P(x=0)$, find the mean of X and P(x s 2).
- 13 (a) Define uniform distribution. Find the moment generating function of the uniform. distribution.
 - (b) In a normal distribution exactly 7% of the items are under 35 and 89% of the items are under 63. Find the mean and standard deviation of the distribution.

14 (a) Find the rank correlation coefficient from the following data:

d the	DOUGH	COURT	CONTRACT	OF E. SON		A 100	24	4.8	22
the x	10	15	12	17.	13.	10	14	1.9	E.E.
2	30	42	45	46	33	34	40	35	39

(b) In two large populations, there are 30% and 25% respectively of fair haired people. Is this difference tikely to be hidden in samples of 1200 and 900 respectively from the two populations? Test at 5% level of significance.

15 Fit a possion distribution to the following data and test the goodness of fit at 5% level of significance.

T.	0	1	2	3	13
f(x)	109	65	22	3	1

18 (a) A continuous random variable X has the probability density

function
$$f(x) = \begin{cases} a + bx, 0 \le x \le 1 \\ 0 & \text{elsewhere.} \end{cases}$$

If the mean of the distribution is $\frac{1}{3}$, find the values of a and b.

- (b) Find the variance of normal distribution.
- 17 (a) If θ is the acute angle between the two regression lines, show that $\tan \theta = \frac{1-r}{r} \cdot \frac{\sigma_s \sigma_s}{\sigma_s^{-1} + \sigma_s^{-1}}$, where $r_r \cdot \sigma_s \cdot \sigma_s$ have their usual meanings. Explain the significance of the formula when r=0 and $r=\pm 1$.
 - (b) Find the least squares approximations of the form y = ar" for the following data:

	0.5				
ď	0.57	1.46	5.10	7.65	9.20

FACULTY OF ENGINEERING B.E. III - Semester (IT) (AICTE) (Main & Backlog) Examination, July 2021

Subject: Mathematics - III (P & S)

Time: 2 Hours

Max. Marks: 70

(Missing data, if any, may be suitably assumed)

PART - A

Note: Answer any five questions.

(5x2 = 10 Marks)

- 1 Let A and B two events such that P(A)=0.5, P(B)=0.6 and P(AUB)=0.8. Find P(A/B).
- 2 If $f(x) = \begin{bmatrix} \frac{k}{n} + k, & 0 \le x \le 3 \\ 0 & \text{otherwise} \end{bmatrix}$ is a probability density function of a random variable, find k.
- 3 Find the moment generating function of binomial distribution.
- 4 A Poisson variant X satisfies P(X=1) = ½ P(X=2). Find the variance of X.
- 5 Find the mean of uniform distribution.
- 6 Write any two properties of normal curve.
- 7. If ==4y+5 and y=k =+4 are two regression lines, show that = 1 = -
- 8 Define level of significance.
- 9 Write the test statistic t to test of significance for difference of means of two small samples.
 10 Write any two uses of F-test.

PART - B

Note: Answer any four questions.

(4x15 = 60 Marks)

- 11 (a) State and prove Baye's theorem.
 - (b) A bag A contains 2 white and 3 red balls and a bag B contains 4 white and 5 red balls. One ball is drawn from one of the bags and is found to be red. Find the probability that it was drawn from bag B.
- 12 (a) If the sum of the mean and variance of a binomial distribution of 5 trials is 9/5, find the binomial distribution.
 - (b) Find the mean and variance of Poisson distribution.
- 13 (a) If a random variable X is uniformly distributed over (-a, a), find 'a' such that P(X>1) =1/3.
 - (b) A continuous random variable X is normally distributed with mean 25 and standard deviation 8. Find the probability that (i) $20 \pm X \pm 40$ and (ii) $|x-25| \le 5$.
- 14 (a) Fit a least square curve of the form y=a, n for the following data:

a:	61	26	7	26
y.	350	400	500	600

(b) A random sample of 900 members has a mean 3.4 cms. Can it be reasonably regarded as a sample from a large population of mean 3.2 cms and standard deviation 2.3 cms? Test at 5% level of significance.

2.

15 The values in two random samples are given below.

Sample 1: 15 25 16 20 22 24 21

Sample II: 35 31 25 38 26 29 32 34 33 27 29 31. Can we conclude that the two samples are drawn from the same population? Test at 5% level of significance.

16 A random variable X has the following probability distribution.

DOMEST WITH	HARMAN NA	11000			Test
1.3	Ti	2	-3	4	0
P(+)	C	C	3c	C*+C	DC.

Find (i) the value of c (ii) E(4X+1)

(iii) Var (4X+1) (iv) P(X<3) and (v) P(1< X <4)

17 Find the correlation coefficient and the equations of regression lines from the following data:

100						-	1.74	T dr	g	10
V	Tr	12	13	4	5	6	1	40	40	50
0	10	12	16	28	25	36	41	491	1	1.4.4

FACULTY OF ENGINEERING

B.E. III Semester (AICTE) (I.T) (Suppl.) Examination, December 2020 Subject: MATHEMATICS-III

Time: 2 Hours

Max. Marks:70

PART - A

Note: Answer any five questions.

(5 x 2 = 10 Marks)

- 1. Write axioms of probability and theorem of total probability.
- State Baye's theorem.
- 3. Find the mean of the Binomial distribution.
- 4. Define skewness of a distribution.
- 5. Explain exponential distribution.
- 6. Find the variance of Uniform distribution.
- Derive the normal equations for a straight line using the principle of least squares.
- Find the angle between the two regression lines.
- 9. Explain level of significance and confidence limits in testing of hypothesis.
- 10. Define Null hypothesis and alternate hypothesis in test of significance.

PART - B

Note: Answer any four questions.

(4 x 15 = 60 Marks)

- 11. a) In a bolt factory, machines */* and c manufacture respectively 25%, 35% and 40% of the total. Of their output 5, 4 and 2 percent are defective bolts. A bolt is drown at random to the product and is found to be defective. What is the probability that it was manufactured by machine if
 - b) is the function defined as follows a density function?

$$f(z) = e^{-z}, z \ge 0$$

= 0 , $z < 0$

If so, determine the probability that variant having this density will fall in the interval (1, 2). Also find the cumulative probability function F(2).

12. a) Fit Poisson distribution to the set of observation

X	0	1	2	3	4
f.	122	60	15	2	1

- b) If the chance that one of the telephone lines is busy at an instant is 0.2
 - i) What is the chance that 5 of the lines are busy?
 - ii) What is the most probable number of busy lines and what is the probability of this number.
 - (iii) What is the probability that all the lines are busy?
- a) A random variable X has Uniform distribution over (-3, 3), find k for which P(X > k)=1/3. Also evaluate P(X < 2) and P(X = 2| = 2).
 - b) In a normal distribution, 7% of the items are under 35 and 89% are under 63.
 What are the mean and S.D of the distribution.

14. a) If P is the pull required to lift a load W by means of pulley block, find a linear law of the from P≈mW+c connecting P and W, using the following data.

P	12	15	21	25
W	50	70	100	120

b) Find the correlation between x and y for the data

x	2	4	6	8	10
y	5	7	9	8	11

- 15. a) The nine items of a sample have the following values 45, 47, 50, 52, 48, 47, 49, 53, 51. Does the mean of these differ significantly from the assumed mean of 47.5?
 - b) The following table gives the number of accidents that took place in an industry during various days of the week. Test if accidents are uniformly distributed over the week.

Day	MON	TUE	WED	THU	FRI	SAT
No of accidents	14	18	12	11	15	14

- 16. a) In a city a 20% of a random sample of 900 school boys had a certain slight physical defect. In another city a , 18.5% of a random sample of 1600 school boys had the same defect. Is the difference between the proportions significant?
 - b) Two random sample from two normal populations are given below:

1	Sample 1	16	26	27	23	24	22
	Sample 2	33	42	35	32	28	31

Do the estimates of population variances differ significantly?

17. a) Find the rank correlation coefficient of the data

7	68	66	68	66	69	66	68	65	71	67	68	70
y	65	63	67	64	68	62	70	66	58	67	69	71

b) In a partially destroyed laboratory record, only the lines of regression of y on a and x on y are available as 20 x - 3 y 10 - 0 and 20 z - 9 y = 107 respectively. Calculate x y and coefficient of correlation between x and y.

Code No. 15050/AICTE

FACULTY OF ENGINEERING

B.E. IV-Samester (EE/Inst/CSE) (AICTE) (Backlog) Examination, March/April 2021

Subject : Mathematics - III (P & S)

Time: 2 hours

Max. Marks: 70

Note: Missing data, if any may be suitably assumed.

PART - A

Answer any five questions.

(5x2 = 10 Marks)

- 1 From a pack of well shuffled cards, one card is drawn. Find the probability that this card is either a king or an ace.
- Check f(x) =
 x = 6 is a probability density function.
- 3 Define akenwess and kurtosis.

4 Find the variance of Binomial distribution.

5 The continuous random variables is uniformly with mean 1 and variance 3. Find p(x < 0).</p>

6 Find the variance of normal distribution.

7 List the properties of correction coefficient.

8 Write the normal equations for fitting the given data as straight line and parabola.

9 Write the test of significance of the difference between the means of two small random samples drawn form the same normal population.

10 Write the test for ratio of variances.

PART - B

Answer any four questions.

(4x15 = 60 Marks)

- 11 (a) A problem in a question paper is given to 3 students in a class to be solved. The probabilities of their solving the problem are 0.5, 0.7 and 0.8 respectively. Find the probability that the problems will be solved.
 - (b) A big contains 5 white and 8 red balls. Two drawing of 3 balls are made such that the balls are not replaced before the second trial. Find the probability that the first drawing ball 3 white and the second 3 red bells in each case.
- 12 (a) An imagular six faced dice is thrown 12 times. The expectation that it will give six even numbers is twice the expectation that it will give 5 even numbers. If 1000 sets, each exactly 12 trials are made, how many sets are expected not to give any even number?
 - (b) In a certain factory manufacturing razor blades, there is a small chance of 0.002 for any blade to be defective. The blades are placed in packets, each containing 10 blades. Using the Poisson distribution, calculate the approximate number of packets containing not more than 2 defective blades in a consignment of 10,000 packets.
- 13 (a) In a distribution which is exactly normal, 12% of the items are under 30 and 85% are under 60. Find the mean and standard deviation of the distribution.
 - (b) Let X₁ and X₂ be two independent uniformly distributed variables in [0, 1]. Then find the distribution of X₁ + X₂

14 (a) Find the correlation coefficient between a and y from the given data :

x	78	89	97	69	59	79	68	57
3	125	137	156	112	107	138	123	108

(b) Fit a parabola y = x + hx + cx to the following data:

8	2	4	- 6	8	10
ÿ.	3.07	12.85	31.47	57 38	01.20

15 (a) A random sample of 17 value from a normal population has a mean of 105 cm and the sum of squares of deviations from this mean is 1225 sq.cm. Is the assumption of 110 cm for the normal population reasonable? Test under 5% and 1% level of significance.

(b) Two random samples of sizes 9 and 7 gave the sum of squares of deviations from their respective means as 175 and 95 respectively. Can they be regarded as drawn from normal

16 (a) Find the best possible curve of the form , -- M, using. Method of least squares for the

7	1	3	4	6.	8	90	34	14
1	1	2	4	4	5	7	8	9

- (b) Two random variables have the regression lines with equations 3x + 2y + 26 and 6x + y = 31. Find the mean values and the correlation coefficient between x and y.
- 17 (a) Fit a Poisson distribution to the following data and test for its goodness of fit at level of

MAN ARIA MAN	_				
No: of defects		1	12	3	1
No of units	214	00	72.65	-	7
	4.77	176	20	3:	1.

(b) The number of students in a class is 100. The average marks scored by 54 boys is 68 with standard deviation of 10 while the average marks scored by 36 girls is 70 with standard deviation of 8. Test of 1% level of significance whether the girls performed better than boys.

FACULTY OF ENGINEERING B.E. III Semester (AICTE) (I.T) (Main)Examination, December 2019

Subject: MATHEMATICS-III

Time: 3 Hours

Max. Marks:70

Note: Answer all questions from Part-A & any five questions from Part-B PART - A (20 Marks)

1. Define Random Experiment, Sample Space, and Event.

2. A Variant X has the Following Distribution.

The second second	STREET, STREET,		
. X	-3	6	9
P(X=x)	1/6	1/2	1/3

Find E (2X+1)2

- 3. Find the Variance of Poisson distribution.
- 4. Define Kurtosis of a distribution.
- 5. Explain Uniform distribution.
- 6. Find the Mean of Exponential distribution.
- Derive the normal equations for the curve y = ax' using the method of least squares.
- Show that correlation coefficient is the geometric mean between the two regression coefficients
- 9. Define Central Limit Theorem.
- 10. Define Type I Error and Type II Error.

PART - B (5 x 10 = 50 Marks)

- 11. a) The content of three ums are, 1 white, 2 rod, 3 green balls, 2 white, 1 red, 1 green balls and 4 white, 5 red, 3 green balls. Two balls are drawn from an um chosen at random. These are found to be one white and one green. Find the probability that the balls so drawn came from the third um.
 - b) The probability density function P(x) of a continuous random variable is P(x) = x,e^{-|x|}, -x < x < x. Prove that x_n = 1/2. Find the mean and variance of the distribution.
- 12. a) Fit a Binomial frequency distribution for the following data.

_						
- X	G	1	2	3	4	5
t	2	14	20	34	22	8

- b) In a certain factory turning out razor blades, there is a small chance of 0.002 for any blade to be defective. The blades are supplied in packets of 10, use Poisson distribution to calculate the approximation number of packets congaing no defectives, one defective and two defective blades respectively in a consignment of 10,000 packets.
- a) Find the variance and moment generating function of uniform distribution. Also find mean, variance the uniform distribution f(x) =1, 0≤ x≤ 1.
 - b) In a Normal distribution, 31% of the items are under 45 and 8% are over 64. Find the mean and standard deviation of the distribution

14 a) An experiment gave the following values

v (ft/min):	350	400	500	600
t (min): (31	26	17	2.6

It is known that v and t are connected by the relation $F=ar^*$. Find the best

b) Find the correlation coefficient between x and y for the following data

				_	-
-	-	2		4	3
3	-1	-	- 9	я	7
y	2	5	3	-	

15. a) The mean of simple samples of size 1000 and 2000 are 67.5cm and 68cm respectively. Can the samples be regarded as drawn from the same population of

b) The theory predicts the proportion of beans in the four groups G1, G2, G3, G4 should be in the ratio 9.3.3.1. In an experiment with 1600 beans the numbers in the four groups were 882, 313, 257 and 118. Does the experiment result support the theory?

16. a) Find the rank correlation coefficient for the following data

	-	-	9	2	7	10	6	6	1	5
20	3	- 8	- 35	4		3.0		4	19	-6
w	5	0	10	- 1	8	173	3	.4		0

b) Two independent sample of sizes 7 and 6 had the following values

Sample A	25	30	32	33	31	29	34
Sample B	29	-30	30	24	27	28	

Examine whether the samples have been drawn from normal populations having the same variance.

17. a) Two random variables have the regression lines with equations 3 x +2 y =26 and 6x + y = 31. Find the mean values and correlation coefficient between x and y

b) Fit the curve y = se" to the following data.

- 1	77	100	185	239	285
	2.4	3.4	7	11.1	19.6

FACULTY OF ENGINEERING B.E. III - Semester (IT) (AJCTE) (Main & Backlog) Examination, July 2021

Subject: Mathematics - III (P & S)

Time: 2 Hours

Max. Marks: 70

(Missing data, if any, may be suitably assumed)

PART - A

Note: Answer any five questions.

(5x2 = 10 Marks)

- 1 Let A and B two events such that P(A)=0.5, P(B)=0.6 and P(AuB)=0.8. Find P(A/B).
- 2 If f(s) = 16 + 3 = 0 = x + 3 is a probability density function of a random variable, find k
- 3 Find the moment generating function of binomial distribution.
- 4 A Poisson variant X satisfies P(X=1) = ½ P(X=2). Find the variance of X.
- 5 Find the mean of uniform distribution.
- 6 Write any two properties of normal curve.
- 7 If x=4y+5 and y=k a +4 are two regression lines, show that a v v
- 8 Define level of significance.
- 9 Write the test statistic t to test of significance for difference of means of two small samples.
- 10 Write any two uses of F-test.

PART - P

Note: Answer any four questions:

(4x15 = 60 Marks)

- 11 (a) State and prove Baye's theorem.
 - (b) A bag A contains 2 white and 3 red balls and a bag B contains 4 white and 5 red balls. One ball is drawn from one of the bags and is found to be red. Find the probability that it was drawn from bag B.
- 12 (a) If the sum of the mean and variance of a binomial distribution of 5 trials is 9/5, find the binomial distribution.
 - (b) Find the mean and variance of Poisson distribution.
- 13 (a) If a random variable X is uniformly distributed over (-a, a), find 'a' such that P(X>1) =1/3.
 - (b) A continuous random variable X is normally distributed with mean 25 and standard deviation it. Find the probability that (i) 20 : X: 40 and (ii) |x - 25| :: 5.
- 14 (a) Fit a least square curve of the form y=a. * for the following data:

40	61	26	7	26
y.	350	400	500	600

(b) A random sample of 900 members has a mean 3.4 cms. Can it be reasonably regarded as a sample from a large population of mean 3.2 cms and standard deviation 2.3 cms? Test at 5% level of significance.

2...

15 The values in two random samples are given below.

Sample I: 15 25 16 20 22 24 21 17 19 23

Sample II: 35 31 25 38 26 29 32 34 33 27 29 31.

Sample II: 35 31 25 38 26 29 32 34 33 27 29 31.

Can we conclude that the two samples are drawn from the same population? Test at 5% level of significance.

16 A random variable X has the following probability distribution.

THE REAL PROPERTY.		and the same of	The state of the s	The same of the sa	
17219	1	2	3	4	5
D/ V	-		3c	C2+C	6c2
	1 1/4	34	- Aprilia		

Find (i) the value of c

(ii) E(4X+1)

(III) Var (4X+1)

(iv) P(X<3) and (v) P(1< X <4).

17 Find the correlation coefficient and the equations of regression lines from the following data:

X	11.	2	3	14	5	6	7	8	9	10
Y	10	12	16	28	25	36	41	497	40	50